

NATIONAL ACADEMY OF SCIENCES

WILMOT HYDE BRADLEY


1899—1979

A Biographical Memoir by
V. E. MCKELVEY

*Any opinions expressed in this memoir are those of the author(s)
and do not necessarily reflect the views of the
National Academy of Sciences.*

Biographical Memoir

COPYRIGHT 1983
NATIONAL ACADEMY OF SCIENCES
WASHINGTON D.C.



W T Bradley

WILMOT HYDE BRADLEY

April 4, 1899–April 12, 1979

BY V. E. MCKELVEY

WILMOT HYDE BRADLEY was both an ordinary and an extraordinary man. He was ordinary in the sense that he was of average build, had plain tastes, was unpretentious, and considered himself to be no better than anyone else. He was, in fact, a superb geologist with extraordinarily broad interests. He was a generalist, not in the sense of one who hasn't specialized in anything or who knows a little bit about a lot of things, but in the sense of one who has demonstrated the ability to probe deeply into diverse subjects and to contribute new and illuminating knowledge about them. He was extraordinary also in his leadership capabilities, exercised first as chief of the Branch of Military Geology, which he helped to found in 1943, and then as chief geologist of the U.S. Geological Survey from 1944 to 1959. In the eyes of his associates he was extraordinary because of his exceptional warmth and selfless attitude toward others, the personal interest he showed in their work and problems, his good humor and wit, and his ability to make almost any conversation an interesting, stimulating exchange among all the participants.

Bill, the son of Anna Miner Hyde and John Lucius Bradley, was born April 4, 1899 in Westville, Connecticut, a suburb of New Haven. He attended grammar school in Westville, high school in New Haven, and college at the Sheffield

Scientific School of Yale University. He enlisted in the U.S. Naval Reserves in 1918, and his junior year at Yale was chiefly given to naval-officer training on a two-masted schooner, supplemented with courses in navigation, spherical trigonometry, and related subjects. Although he had majored in engineering in his first years at Yale, he switched to chemistry in his senior year. He soon had doubts about his choice, however, and his friend and later USGS colleague, Arthur A. Baker, then a first-year graduate student, suggested that geology might be more to his liking. After nine weeks of exposure to an introductory course taught by Alan Bateman, he changed his major to geology and graduated from Yale in 1920 with a Ph.B. That summer he served as field assistant to Frank C. Calkins of the U.S. Geological Survey in the Cottonwood District in the Wasatch Mountains of Utah.

The following two years were spent in graduate studies at Yale, with summers as a geologic aide to Julian D. Sears of the Survey on the north flank of the Uinta Mountains. During the second of these field seasons with "J.D.," in which James G. Gilluly also served as an assistant, Bill first saw and became fascinated with the Eocene Green River Formation. He learned that David White, then chief geologist, was looking for someone to work full time on the Green River because of its oil shale potential; he volunteered for the assignment and was taken on full time by the Survey in the fall of 1922 to work on the Green River.

The fall of 1922 thus began Bill's union with two of his lifelong loves—the Green River Formation and the Survey. A third was joined during the same period when, on November 4, he married Catrina van Benschoten, also of New Haven and a friend since childhood. Their marriage—blessed with two daughters, Anne and Penny—was a devoted one, lasting until Bill's death.

Bill's Geological Survey Professional Paper 140, "Shore Phases of the Green River Formation in Northern Sweet-water County, Wyoming," served also as his doctoral dissertation, and he received his Ph.D. from Yale in 1927. His sound academic training laid a solid foundation for the career that followed. He was appreciative of his teachers' stimulus—particularly Adolph Knopf at Yale who inspired him to search for causes and dependent relationships among natural phenomena and processes.

Bill's broad interests had been stimulated before he reached the college level by his father, a dentist, who was interested in all things mechanical and electrical, and who taught Bill how to wire motors and to make and experiment with various kinds of wet batteries, Leyden jars, and other electrical devices. His mother and her maiden sister, Carolyn, also played strong parts in arousing Bill's curiosities, for they were intensely interested in birds, flowers, butterflies, and moths and took him on numerous trips to nearby woods and meadows, and to Yale University's Peabody Museum as well.

With this background, it is easy to understand how Bill's interest in the Green River Formation encompassed almost every aspect of its composition and geologic origin. His first scientific paper described "Fossil Caddice Fly Cases from the Green River Formation of Wyoming," and subsequent papers dealt with its mineralogy, plant and animal fossils, physical structures such as varves and mud cracks, stratigraphy and areal geology, and geochemistry, as well as the climate of the Green River Epoch and the paleolimnology of the Green River lakes. In his later years, Bill broadened his study of oil shale to its formation in the modern environment. Motoaki Sato, Bill's colleague in some of these studies, says of them:

One of Bradley's main scientific concerns was to find a modern lake that was producing rich organic ooze with very little clastic material. He was

excited when he heard R. S. A. Beauchamp, an English limnologist, describe a remarkable organic ooze that was forming in the northern part of Lake Victoria. The organic ooze consisted almost wholly of algal matter which would not decay in a warm, wet, and oxidizing environment. Bradley immediately began his search for such lakes and found that only one more lake in equatorial East Africa and two lakes in Florida, one of which is Mud Lake, are known to be accumulating this kind of pure algal ooze. His tireless and all-out effort in understanding the limnology, microbiology, and geochemistry of Mud Lake, Marion County, Florida, began soon afterwards. Bradley's approach was characteristically multidisciplinary. Not only did he examine the algal ooze microscopically to identify microorganisms and the evidence for their activities, he mobilized experts in the nation to identify various organic compounds and microbes existing in the organic sediment, and to develop tools for sampling the ooze and conducting *in situ* measurements of geochemical parameters. His main effort was directed to unravelling the secret of how the algal matter resisted decay in a subtropical to tropical environment and how relatively oxygen-rich algal matter changed with time to the hydrocarbon-rich organic matter of oil shale in an ordinary diagenetic environment. His pioneering efforts in this realm of science have given impetus to many students of organic sedimentation. One of the most thorough documentations of geochemical parameters existing in organic sediments, which Bradley worked on even after his retirement from the U.S. Geological Survey, is expected to be published shortly.¹

With respect to the significance of Bill's overall work on the Green River, Erle Kauffman of the U.S. National Museum writes:

Bill Bradley was the true "father" of non-marine aquatic paleoecology and paleolimnology. He pioneered modern technique by crossing the line between recent and ancient ecosystems to become a respected limnologist and aquatic biologist, and then to apply that knowledge with great precision to the interpretation of ancient ecosystems. His early work was twenty to thirty years ahead of its time, and stands today as one of the best examples of fresh water paleoecology on record. A measure of Bradley's perception as a student of paleoecology and paleoenvironments has come to light in the recent "Green River controversy," which seemed to pit

¹Motoaki Sato 1979: personal communication.

Bradley's older stratified fresh water lake model against a newer playa lake model for the origin of Green River oil shale. Bradley reasoned that varved oil shales with their high organic residues and beautifully preserved biota could only have formed in a permanently stratified lake, characterized by a thick, poorly circulated, O₂ depleted and H₂S enriched hypolimnion and a metalimnion and epilimnion with a diverse fresh water biota, high productivity, and seasonal fluctuation producing well defined varves.

From the outset, Bill recognized a significant stratigraphic interruption in the middle of the Green River formation in which the stratified lake model broke down. He termed this the "middle saline facies," or the Wilkins Peak Member of the Green River Formation. He noted paleogeographic restriction of the Wilkins Peak and determined that the lake had shrunk considerably and was without outlet at this time. He reviewed the geochemistry of the unit, and especially the unique suite of authigenic saline minerals. From this he concluded that evaporation greatly exceeded fresh water input, and that a shallow, clear saline lake persisted, with broad episodically exposed brine flats. Bradley was describing a playa; he simply never called it this. Subsequent filling of the Green River lake basins resulted in the redevelopment of a large stratified freshwater lake system, possibly with a saline hypolimnion. Subsequent detailed stratigraphic analyses of the Wilkins Peak Member by others have revealed extensive new evidence in support of the playa lake model for the Wilkins Peak phase of the Green River. Whereas this started out as a sober analysis of a specific unit, it mushroomed into a quiet controversy in which some workers insisted on a playa origin for most of the Green River System. From the outset, Bradley agreed that the Wilkins Peak evidence fit the playa model, and even presented new evidence in support of it. But at the same time he quietly warned of extrapolating to make one depositional system (the playa model) fit the whole Green River. His prediction was correct, the controversy has run its course, and now in the aftermath Bradley's perception and the breadth of his observations have re-emerged. His original stratified lake model, from the standpoint of integrated geochemical, sedimentological, paleogeographic, and paleoecological evidence, is still supported for most of Green River time, and most of the varved oil shale.²

Although the Green River Formation was the stimulus for much of Bill's work, his field studies included geologic map-

² Erle Kauffman 1979: personal communication.

ping and evaluation of oil and gas possibilities in southwestern Wyoming and adjacent parts of Utah and Colorado, as well as in southcentral New York. One of the most rewarding of his other assignments was his pioneering work on the C. S. Piggott cores from the deep ocean floor of the North Atlantic between the Grand Banks off Newfoundland and the continental shelf of Ireland. Eleven in all, and averaging nearly 8 feet in length, they were the first obtained from the abyssal depths. Bill's careful studies of these cores, made in collaboration with several others, showed for the first time the possibilities of unravelling geologic, oceanographic, and climatic history from analysis of the sedimentary record on and beneath the deep ocean floor.

Another series of investigations demonstrating Bill's multidisciplinary interests had to do with the dynamics and history of tidal flats in Maine. It is noteworthy that his principal report on these studies was published by the U.S. Fish and Wildlife Service because of their bearing on commercial clamming.

Although the Geologic Division of the Geological Survey had been blessed with fine leadership beginning with its first chief geologist, Grove Karl Gilbert, none gave it better or more dedicated service than Bill Bradley. Wise in the ways in which a scientific organization can be guided to do its best work, Bill led not by command or directive; through inspiration, the contagiousness of his own enthusiasm, and the stimulating effect of his interest in other people and their work and problems, he brought forth their best efforts. Bill's sixteen-year period of service as a branch chief and chief geologist spanned a period of great change in the Geologic Division, including a several-fold expansion in professional personnel and the beginning or expansion of important new programs, such as military geology, airborne geophysics, the geology of radioactive minerals, and engineering geology.

Not only was he able to find solutions to the many difficult problems that he had to face during this expansion, but he found them without affronting or offending those concerned. He won and held the deep respect and affection of his associates.

Through his membership, Bill contributed to the activities of a diverse array of organizations: National Academy of Sciences, American Philosophical Society, American Academy of Arts and Sciences, Geological Society of America, American Society of Limnology and Oceanography, International Limnological Association, American Association for the Advancement of Science, American Association of Petroleum Geologists, Botanical Society of America, Sigma Xi, Society of Economic Paleontologists and Mineralogists, Geological Society of Washington, and the Cosmos Club of Washington. Several of these organizations honored him with awards and high office: the National Academy with its Award of Merit in 1940, the Philadelphia Academy of Science with the F. V. Hayden Medal and Award in 1971, the Geological Society of America with its presidency in 1965 and its Penrose Medal in 1972, the Geological Society of Washington with its presidency in 1946, and the Society of Economic Paleontologists and Mineralogists with honorary membership. Bill also received the Department of the Interior's Distinguished Service Award in 1958 and an honorary Doctor of Science degree from Yale in 1947. His colleagues honored him with the *Bradley Volume Festschrift*, published by the *American Journal of Science* in 1960.

At the conclusion of his forty-eight-year career with the Geological Survey in 1970, Bill and Catrina moved to the west shore of Pigeon Hill Bay, Maine. There he continued writing up his results on the Green River and Mud Lake while enjoying the physical stimulation of outdoor work on their farm. During a visit with him in the fall of 1978, when illness had

already begun to overtake him, he was rhapsodic about the enjoyment and satisfaction his life had given him—the love of his family, the excitement of his research, his stimulating and rewarding friendships, and his life with the Geological Survey. His life was a joyous and satisfying one to him and an enriching one for his family, his friends, his scientific organizations, his country and its Geological Survey, and his science. He died of a stroke on April 12, 1979, eight days after his eightieth birthday.

BIBLIOGRAPHY

1924

- Fossil caddice fly cases from the Green River Formation of Wyoming. *Am. J. Sci.*, 5th ser. 7:310-12.
- An oil shale and its microorganisms from the Fuson Formation of Wyoming. *Am. J. Sci.*, 5th ser. 8:228-34.

1925

- A contribution to the origin of the Green River Formation and its oil shale. *Am. Assoc. Pet. Geol. Bull.*, 9(2):247-62.

1926

- Shore phases of the Green River Formation in northern Sweet-water County, Wyoming. *U.S. Geol. Surv. Prof. Pap.*, 140: 121-31.

1928

- Zeolite beds in the Green River Formation. *Science*, n.s. 67:73-74.

1929

- The occurrence and origin of analcite and meerschaum beds in the Green River Formation of Utah, Colorado, and Wyoming. *U.S. Geol. Surv. Prof. Pap.*, 158:1-7.
- The varves and climate of the Green River epoch. *U.S. Geol. Surv. Prof. Pap.*, 158:87-110.
- Algae reefs and oolites of the Green River Formation. *U.S. Geol. Surv. Prof. Pap.*, 154:203-23.
- Cultures of algal oolites. *Am. J. Sci.*, 5th ser. 18:145-48.
- Fresh-water algae from the Green River Formation of Colorado. *Torrey Bot. Club Bull.*, 56(8):421-28.
- Neue Beobachtungen über Algen als Urmaterialien der Bogheadkohlen und schiefer. *Centralbl. Min., Jahrg., Abt. B. Nr. 5.S.*:182-90.

1930

- The behavior of certain mud-crack casts during compaction. *Am. J. Sci.*, 5th ser. 20:136-44.

1931

Origin and microfossils of the oil shale of the Green River Formation of Colorado and Utah. U.S. Geol. Surv. Prof. Pap. 168. 58 pp.

Nonglacial marine varves. *Am. J. Sci.*, 5th ser. 22:318-30.

1933

Factors that determine the curvature of mud-cracked layers. *Am. J. Sci.*, 5th ser. 26(151):55-71.

1935

Geology of the Alcova Dam and reservoir sites, North Platte River, Natrona County, Wyoming. *Econ. Geol.*, 30(2):147-65.

Anticlines between Hiawatha gas field and Baggs, Wyoming. *Am. Assoc. Pet. Geol. Bull.*, 19(4):537-43.

Structure and gas possibilities of the Watkins quadrangle, New York. U.S. Dept. Interior Press Memo 101944. 14 pp.

1936

Geomorphology of the north flank of the Uinta Mountains (Utah). U.S. Geol. Surv. Prof. Pap. 185-I:163-99.

1937

The biography of an ancient American lake. *Sci. Mon.*, 42(5): 421-30. (Reprinted, *Smithson. Inst. Annu. Rep.*, 1937.)

Nonglacial varves, with selected bibliography. *Natl. Res. Council. Annu. Rep. Appendix A*:32-42.

With M. N. Bramlette, J. A. Cushman, L. G. Henbest, K. E. Lohman, and P. D. Trask. Preliminary report on the North Atlantic deep-sea cores taken by the Geophysical Laboratory, Carnegie Institution. *Am. Geophys. Union Trans.*, 18th Annu. Mtg., part 1:224-26.

1938

With J. F. Pepper. Structure and gas possibilities of the Oriskany sandstone in Steuben, Yates, and parts of the adjacent counties, New York. U.S. Geol. Surv. Bull. 889-A. 68 pp.

With M. N. Bramlette, J. A. Cushman, L. G. Henbest, K. E. Lohman, and P. D. Trask. The Geological Survey's work on the Piggott North Atlantic deep sea cores. *Proc. Am. Philos. Soc.*, 79(1):41-46.

A brief annotated bibliography on cyclic variations in climate as indicated by pre-Pleistocene nonglacial varves. *Bull. Am. Meteorol. Soc.*, 19(5):162-63.

Mediterranean sediments and Pleistocene sea levels. *Science*, 88(2286):376-79.

1940

Geology and climatology from the ocean abyss. *Sci. Mon.*, 50(2): 97-109.

Pediments and pedestals in miniature. *J. Geomorphol.*, 3(3): 244-54.

1942

With M. N. Bramlette. Geology and biology of North Atlantic deep-sea cores between Newfoundland and Ireland. *U.S. Geol. Surv. Prof. Pap.* 196. 157 pp.

1945

Geology of the Washakie Basin, Sweetwater and Carbon Counties, Wyoming, and Moffat County, Colorado. *U.S. Geol. Surv. Oil Gas Invest. Prelim. Map* 32.

1946

Coprolites from the Bridger Formation of Wyoming, their composition and microorganisms. *Am. J. Sci.*, 244(3):215-39.

1947

A suggested geological curriculum. *Geol. Soc. Am. Interim Proc.*, part 2:8-13.

1948

Limnology and the Eocene lakes of the Rocky Mountain region. *Geol. Soc. Am. Bull.*, 59(7):635-48.

1953

Age of intertidal tree stumps at Robinhood, Maine. *Am. J. Sci.*, 251:543-46.

1957

Radiocarbon age of Damariscotta shell heaps. *Am. Antiq.*, 23:3.
Physical and ecologic features of the Sagadahoc Bay tidal flat, Georgetown, Maine. *Geol. Soc. Am. Mem.* 67:641-82.

1959

With Peter Cooke. Living and ancient populations of the clam *Gemma gemma* in a Maine coast tidal flat. U.S. Fish Wildl. Serv. Fish. Bull., 58(137):305-55.

Revision of the stratigraphic nomenclature of the Green River Formation of Wyoming. *Am. Assoc. Pet. Geol. Bull.*, 43(5):1072-75.

1961

Geologic map of a part of southwestern Wyoming and adjacent states. U.S. Geol. Surv. Misc. Geol. Invest. Map I-332.

1962

With J. J. Fahey. Occurrence of stevensite in the Green River Formation of Wyoming. *Am. Mineral.*, 47:996-98.

Chloroplast in *Spirogyra* from the Green River Formation of Wyoming. *Am. J. Sci.*, 260:455-59.

Memorial to Esper Signius Larsen, 3d. *Geol. Soc. Am. Proc.*:35-37.

1963

Continental Divide split. *Geotimes*, 8(3):26.

Unmineralized fossil bacteria. *Science*, 141:919-21.

Geologic laws. In: *The Fabric of Geology*, ed. Claude C. Albritton, pp. 12-23. Reading, Mass.: Addison-Wesley.

Paleolimnology. In: *Limnology in North America*, ed. David G. Frey, pp. 621-52. Madison: University of Wisconsin Press.

1964

Aquatic fungi from the Green River Formation of Wyoming. *Am. J. Sci.*, 262:413-16.

Lazurite, talc, and chlorite in the Green River Formation of Wyoming. *Am. Mineral.*, 49(5-6):778-81.

Geology of Green River Formation and associated Eocene rocks in southwestern Wyoming and adjacent parts of Colorado and Utah. U.S. Geol. Surv. Prof. Pap. 496-A. 86 pp.

1965

Vertical density currents. *Science*, 150:1423-28.

1966

Paleolimnology of the trona beds in the Green River Formation of Wyoming. In: *Second Symposium on Salt*, pp. 160-64. Cleveland: Northern Ohio Geological Society.

Memorial to Levi Fatzinger Noble. *Geol. Soc. Am. Bull.*, 77: P49-P52.

Tropical lakes, copropel, and oil shale. *Geol. Soc. Am. Bull.*, 77: 1333-38.

1967

Two aquatic fungi (Chytridiales) of Eocene age from the Green River Formation of Wyoming. *Am. J. Bot.*, 54(5) part 1:577-82.

Precursors of oil shale. *Proceedings of Seventh World Petroleum Congress, Mexico City, April 1967*, pp. 695-97. Great Yarmouth, England: Galliard.

1968

Unmineralized fossil bacteria: a retraction. *Science*, 160:437.

1969

Vertical density currents—II. *Limnol. Oceanog.*, 14(1):1-3.

Walter Herman Bucher. In: *Biographical Memoirs*, 40:19-34. New York: Columbia University Press for the National Academy of Sciences.

With H. P. Eugster. Geochemistry and paleolimnology of the trona deposits and associated authigenic minerals of the Green River Formation of Wyoming. U.S. Geol. Surv. Prof. Pap. 496-B. 69 pp.

With K. B. Hoag, A. J. Tousimis, and D. L. Price. A bacterium capable of using phytol as its sole carbon source, isolated from algal sediment of Mud Lake, Florida. *Proc. Natl. Acad. Sci. USA*, 63(3):748–52.

Memorial to Carle Hamilton Dane. *Geol. Soc. Am. Proc.* 1968:1–7.

1970

With M. E. Beard. Mud Lake, Florida; its algae and alkaline brown water. *Limnol. Oceanog.*, 14:889–97.

With A. J. Iovino. The role of larval Chironomidae in the production of lacustrine copropel in Mud Lake, Marion County, Florida. *Limnol. Oceanog.*, 14:898–904.

Green River oil shale—concept of origin extended. *Geol. Soc. Am. Bull.*, 81(4):985–1000.

Eocene algae and plant hairs from the Green River Formation of Wyoming. *Am. J. Bot.*, 57(7):782–85.

1973

Oil shale formed in desert environment: Green River Formation, Wyoming. *Geol. Soc. Am. Bull.*, 84:1121–24.

1974

Oocardium tufa from the Eocene Green River Formation of Wyoming. *J. Paleontol.*, 48:1289–90.

In Press

With M. Sato. Electrochemical probing in sulfide-rich environments having steep temperature gradients. *Am. J. Sci.*

With M. Sato. Vertical variation of geochemical environment of algal sediment in Mud Lake, Florida. *Am. J. Sci.*